



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Distortion product otoacoustic emission fine structure of symphony orchestra musicians

Reuter, Karen; Hammershøi, Dorte

Publication date:
2006

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Reuter, K., & Hammershøi, D. (2006). *Distortion product otoacoustic emission fine structure of symphony orchestra musicians*. Poster presented at Inter-Noise 2006, Honolulu, Hawaii, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.



Distortion Product Otoacoustic emission fine structure of symphony orchestra musicians

Karen Reuter and Dorte Hammershøj

Acoustics, Department of Electronic Systems, Aalborg University, Denmark

Introduction



In this study distortion product otoacoustic emission (DPOAE) were measured with a high frequency resolution for a group of musicians before and after rehearsal. The high frequency resolution reveals a fine structure [1-11], which is the result of two or more distortion product sources on the basilar membrane. The primary component comes from a place close to f_2 , and the secondary (reflection component) from the $2f_1-f_2$ site. The fine structure will be more or less pronounced, depending on the strength relations between the two components, which again depends on the state of hearing (e. g. OHC damage).

The purpose was to study any temporary effect from rehearsal, and to study any preliminary permanent effect from work-conditioned exposure. For details, see [12].

Methods

$2f_1-f_2$ DPOAE measurements were done with the ILO96: $L_1/L_2 = 65/45$ dB, $f_2/f_1 = 1.22$, 903 Hz $> f_2 > 6201$ Hz, $\Delta f = 12$ Hz for $f_2 < 3$ kHz and $\Delta f = 22$ Hz for $f_2 > 3$ kHz. 12 musicians aged 31-55 (mean 39) participated, Hearing thresholds were measured using standard audiometry.

One subject participated per rehearsal day. Each session lasted 90 minutes, 30 minutes for the pure-tone audiometry and 60 minutes for the measurement of DPOAE fine structure. The order of testing either hearing threshold or DPOAE first was balanced.




DPOAE measurement for one subject before and after rehearsal. The example features fine structure characteristics with deep notches and high peaks.


DPOAE mean of 12 subjects before and after rehearsal. The individual fine structures are smoothed out, when the data are averaged across subjects.

There was no significant DPOAE level shift after the rehearsal, nor threshold shift. This may be partly due to the fact that some time elapsed from cessation of the rehearsal until the measurements could begin. Another explanation is that the measurements took quite some time, and the hearing therefore recovered before and/or during measurements.

The data from the musicians were compared with an age and sex balanced control group, and further compared with data for young normal-hearing students subdivided into the best and slightly elevated thresholds (above 10 dB HL), from [13].



Musicians and the control group compare well, and have relatively good hearing for their age. The best hearing younger individuals (best-hearing students) have thresholds that compare well to the reference.




DPOAE data are also clearly better for best-hearing subjects. Musicians and control group compare well.


DPOAE fine structure

Fine structure characteristics were determined using the algorithm presented in [1], which determines:


- maxima and minima of fine structure ripples,
- ripple spacing as frequency difference between two minima,
- ripple height as level difference between maxima and mean of the two minima,
- number of ripples as ripples > 3 dB in height per $1/3$ octave.



The ripple spacing is quite similar for the four groups, although on average slightly more narrow for the group of students with slightly elevated thresholds.



The ripple height is also quite similar for the four groups, although on average slightly higher for the group of students with slightly elevated thresholds.



The ripple prevalence is also quite similar for the four groups, although with a distinct bend around 3 kHz for the group of students with slightly elevated thresholds.

Conclusion

There was no difference in the state of hearing for the musicians before and after rehearsal. The comparison between musicians and the control group, which were sex- and age-balanced to the musicians, revealed no significant difference in any measure. Both groups differed from the best-hearing students, both in overall DPOAE level and by some of the fine structure characteristics (spacing and height).

The difference in fine structure characteristics has no simple explanation. It may be that the early age-related deterioration of the hearing leads to more irregularities on the basilar membrane. These may generally affect the secondary (reflection) component for more frequencies than the place-specific component, thus the higher number of dips is related to higher order reflections.

The higher number of dips may however also be a consequence of the lower overall DPOAE level, which may result in a higher number of false dip detections.

Acknowledgements

Financial support of the William Demant Foundation (Oticon) and the Danish Research Council for Technology and Production is sincerely acknowledged. The cooperation with Aalborg Symphony Orchestra is greatly acknowledged. Thanks to all musicians and other individuals for participating in this experiment, and to Rodrigo Ordoñez for assistance.

References

1. Gaskill and Brown (1990) J. Acoust. Soc. Am. 88:821-839.
2. He and Schmidt (1993) J. Acoust. Soc. Am. 94:2659-2669.
3. Heitmann et al. (1996) Eur. Arch. Otorhinolaryngol. 253:167-171.
4. Mauermann et al. (1999) J. Acoust. Soc. Am. 106:3473-3483.
5. Mauermann et al. (1999) J. Acoust. Soc. Am. 106:3484-3491.
6. Talmadge et al. (1998) J. Acoust. Soc. Am. 104:1517-1543.
7. Talmadge et al. (1999) J. Acoust. Soc. Am. 105:275-292.
8. Knight and Kemp (2000) J. Acoust. Soc. Am. 107:457-473.
9. Kalluri and Shera (2001) J. Acoust. Soc. Am. 109:622-637.
10. Konrad-Martin et al. (2002) J. Acoust. Soc. Am. 111:1800-1809.
11. Long et al (2004) Abstract, ARO mid winter meeting.
12. Reuter and Hammershøj (2007) J. Acoust. Soc. Am. 121:xx-xx.
13. Reuter and Hammershøj (2006) J. Acoust. Soc. Am. 120:270-279.